

ACS/STEMI KPI Metric				
Metric	Definition	Benchmark	Numerator/Denominator	
FMC to EKG	Elapsed time, in minutes, from first medical contact to initial 12 lead EKG acquisition	<= 8 min, 90% of cases where ACS/STEMI is suspected		
Stroke KPI Metric				
Metric	Definition	Benchmark	Numerator/Denominator	
Transport initiated within 15 minutes	Begin transport within fifteen minutes of arrival of the first EMS unit on scene	85% for Stroke patients		
BGL	Documentation of a blood glucose level obtained via capillary sample or the recommended sample per manufacture's instructors for the device	Obtained and documented, 100% of all patients with suspected stroke		
Intubation KPI Metric				
Metric	Definition	Benchmark	Numerator/Denominator	
First pass success on endotracheal intubation	First attempt at intubation success. Attempt is defined as placement of the laryngoscope into the mouth	75% First pass success for all Intubations		
Supraglottic success after missed intubation	Failed ET attempt converted to the use of a supraglottic airway after the first ET attempt	Total number of conversions		
Weights KPI Metric				
Metric	Definition	Benchmark	Numerator/Denominator	
Documentation of pediatric patient weight on PCR	All pediatric patients should have a weight documented on the EPCR to ensure accurate medication dosages used	Enter weight (in kg) on ePCR of at least 95% of patients between 0-18 years of age		
Pain Management KPI Metric				
Metric	Definition	Benchmark	Numerator/Denominator	
Documentation of a pain scale pre & post treatment	All patients that receive pharmacological pain treatment or other non-invasive treatment (splints, position of comfort, ice, etc.) documented on the EPCR	Enter scale on ePCR of at least 95% for patients		
BLS Specific KPI Metric				
Metric	Definition	Benchmark	Numerator/Denominator	
Temperature determination of suspected sepsis patients	Documented assessment of subjective (e.g. tactile) or objective (e.g. measured) temperature	85% of cases for patients with suspected or confirmed infection		
Documented Blood Glucose on AMS patients	All patients with any altered mental status should have BGL taken and recorded on the EPCR	90% of BLS AMS patients have an initial BGL taken		
Documentation of pain scale on all patients with pain receiving EMS treatment	All patients that receive pharmacological pain treatment or other non-invasive treatment (splints, position of comfort, ice, etc.) documented on the EPCR	Enter scale on ePCR of at least 95% for patients		